ISSN:2583-9071
VOLUME 3, ISSUE 2 I FEB 2025

A STEP TOWARDS AGRICULTURE

Agri Roots

e-Magazine

Union Budget 2025: What to Expect for Agriculture

FEBRUARY 2025
WWW.AGRIROOTSMAGAZINE.IN

FARMER WELFARE, BOOSTING RURAL INFRASTRUCTURE, AND INVESTING IN AGRI-TECH. THESE CAN EMPOWER THE BACKBONE OF OUR ECONOMY.

TABLE OF CONTENT

MEET *the* EDITORIAL BOARD

Dr. Deepak Kumar
Founder & Editor
Assistant Professor, School of Agriculture, Shobhit Deemed to-be University, Meerut (Uttar Pradesh), India

Dr. Vipin Kumar
Editor-in-Chief
Professor, College of Horticulture, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut (Uttar Pradesh), India

Dr. B.S. Tomar
Editor
Head, Professor And Principal Scientist Division Of Vegetable Science, ICAR - Indian Agricultural Research Institute, New Delhi, India

Dr. Moolchand Singh
Editor
Principal Scientist, Division Of Plant Quarantine ICAR-NBPGR (Ministry Of Agriculture & Farmers Welfare, Govt. Of India) IARI Campus (New Delhi), India

Dr. Ravindra Kumar

Editor

Senior Scientist, Plant Pathology, Crop Protection Division, ICAR- Indian Institute Of Wheat And Barley Research, Karnal (Haryana), India

Dr. Dilpreet Talwar

Editor

Extension Scientist (Vegetable), College Of Horticulture And Forestry, Punjab Agricultural University, Ludhiana (Punjab) India

Dr. Pankaj Kumar Kannaujia

Editor

Scientist (Senior Scale), Horticulture, Division Of Plant Exploration And Germplasm Collection, ICAR- NBPGR (New Delhi), India

Dr. Vineet Kumar

Editor

Assistant Professor, Department Of Soil Science, Galgotias University, Greater Noida (Uttar Pradesh) , India

MEET *the* REVIEWER BOARD

Dr. Khushboo Kathayat
Reviewer
Assistant Professor, Department of Horticulture, College of Agriculture. Lovely Professional University, Phagwara (Punjab), India

Dr. Pratima Gupta
Reviewer
Subject Matter Specialist, KVK, Nagina, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (Uttar Pradesh), India

Dr. Sauhard D
Reviewer
Subject Matter Specialist, KVK, Ujhani, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (Uttar Pradesh), India

Dr. Pankaj Kumar
Reviewer
Subject Matter Specialist, KVK, Chandausi, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut (Uttar Pradesh), India

Dr. Aman Deep Ranga
Reviewer
Ph.D., Dr. YS Parmar University of Horticulture and Forestry, Solan, Himachal Pradesh, India

Dr. Sudhir Kumar
Reviewer
Assistant Professor, Department of Food Technology, School of Advanced Agriculture Sciences & Technology, CSJMU, Kanpur

Dr. Gaurav Tripathi
Reviewer
PDF, Geo-Informatics, Department of Civil Engineering, Indian Institute of Technology, Bombay (India)

Mr. Navdeep Singh
Reviewer
Ph.D. Scholar, Horticulture (Vegetable Science), Punjab Agricultural University, Ludhiana (Punjab), India

from the editor

As the Indian government prepares for the Union Budget of 2025, expectations are high that it will address the pressing challenges faced by the agricultural sector. Agriculture, which continues to employ a large portion of the Indian workforce, is grappling with issues ranging from stagnating productivity and environmental stress to the ongoing struggle with farmer income and inadequate infrastructure.

One of the key areas of focus in the upcoming budget is expected to be a renewed push for farmers' welfare. It is anticipated that the government will allocate more funds for direct income support schemes, such as the PM-KISAN program, to ensure a sustainable livelihood for farmers, particularly smallholders who often remain vulnerable to price volatility and climate shocks.

The sustainability of farming practices is likely to gain further attention, especially in the context of climate change. We may see increased investment in green technologies, such as precision farming, water management systems, and the promotion of organic farming. Given the expanding interest in agricultural export markets, it is expected that the budget will emphasize enhancing agricultural trade infrastructure. Initiatives aimed at improving logistics, building export hubs, and facilitating faster movement of goods could open new opportunities for Indian farmers in global markets. Additionally, the creation of a more efficient agri-tech ecosystem is likely to be a major focus, with provisions for digital platforms and data-driven decision-making tools that can empower farmers and improve productivity.

Dr. Deepak Kumar
FOUNDER & EDITOR

EXPLORING KNOWLEDGE & DISCOVERING AGRICULTURE

AGRI ROOTS E-MAGAZINE

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

New Education Policy in Agricultural Education

ARTICLE ID: 0177

Dinesh Kumar Meena

[1]Department of Horticulture, School of Agricultural Sciences and Technology Babasaheb Bhimrao Ambedkar University (A Central University) Vidya Vihar, Raebareli Road, Lucknow-226025, (UP.)- India

The New Education Policy (NEP) of India aims to transform the educational landscape, ensuring inclusivity, accessibility, and quality across disciplines, including agricultural education. This chapter delves into the alignment of NEP with agricultural education, highlighting its potential to address sustainable development goals (SDGs), social equity, and agricultural innovation. It underscores the importance of reimagining agricultural curricula, fostering interdisciplinary approaches, and integrating modern technology to prepare future professionals for emerging challenges in the agricultural sector.

Agriculture is the backbone of India's economy, employing nearly half of the workforce and contributing significantly to GDP. With increasing population, climate change, and resource constraints, the agricultural sector faces numerous challenges that demand innovative solutions. Education plays a pivotal role in equipping individuals with the knowledge and skills required to address these issues. The New Education Policy (NEP) 2020 provides a comprehensive framework to revamp the Indian education system, including agricultural education, aligning it with contemporary needs and global standards.

New Education Policy

The New Education Policy 2020 is a progressive reform introduced by the Government of India to overhaul the education system. It emphasizes holistic, multidisciplinary, and skill-based learning, aiming to create lifelong learners. Key pillars of NEP include:

- Flexibility in curriculum and pedagogy.
- Focus on vocational and skill development.
- Promotion of research and innovation.
- Integration of technology in education.
- Inclusion and equity.

Integration of NEP Principles in Agricultural Education

NEP Principle	Implementation in Agricultural Education	Innovative Example	Impact
Multidisciplinary Learning	Merging agriculture with technology, management, and arts	Agri-tech courses, agri-entrepreneurship	Holistic skill development
Flexibility in Curriculum	Choice-based credit system (CBCS)	Students opting for agribusiness or organic farming modules	Customizable learning paths
Skill-based Education	Focus on practical and vocational skills	Workshops on drone usage, hydroponics	Industry readiness
Research & Innovation	Promoting problem-solving through real-world projects	Field labs for climate-resilient crops	Enhanced problem-solving abilities
Digital and Online Learning	Blending traditional teaching with online platforms	MOOCs in sustainable farming practices	Wider access and updated knowledge

NEP-driven Curriculum Innovations in Agricultural Education

Curriculum Component	NEP Feature Integrated	Example	Expected Outcome
Multidisciplinary Projects	Collaboration with other disciplines	Joint courses with engineering and business	Cross-functional expertise
Internships and Apprentices	Vocational focus, industry exposure	Internship at agribusiness firms	Real-world experience
Entrepreneurship Modules	Fostering innovation and startups	Agri-entrepreneurship workshops	Startup culture in rural areas
Blended Learning	Online and offline integration	Online classes on emerging technologies	Greater accessibility to rural students
Global Exposure	International collaborations	Exchange programs with global institutes	Exposure to global best practices

Importance of New Education Policy

The NEP aims to address systemic gaps and build a robust educational framework to meet global challenges. Its importance in agricultural education includes:

1. Modernizing curricula to incorporate emerging technologies like AI, IoT, and precision farming.
2. Promoting interdisciplinary learning to integrate agricultural sciences with environmental studies, economics, and management.
3. Emphasizing research and entrepreneurship to foster innovation in agriculture.
4. Encouraging practical, hands-on training to prepare students for real-world challenges.

Interlink of New Education Policy and Agricultural Education

The NEP's provisions significantly impact agricultural education:

1. **Holistic and Multidisciplinary Learning**: The NEP promotes a multidisciplinary approach, enabling students to blend agriculture with allied fields like biotechnology, data science, and environmental management.
2. **Vocational Training**: Strengthened focus on vocational training ensures practical exposure to modern farming techniques and agribusiness.
3. **Digital and Technology Integration**: The NEP encourages leveraging digital platforms for remote learning, enabling wider access to quality agricultural education.
4. **Innovation and Research**: Establishing research-intensive institutions fosters cutting-edge developments in sustainable agriculture and food security.

NEP, SDGs, and Agricultural Education

Agriculture is central to achieving several SDGs, including zero hunger, sustainable cities, and climate action. The NEP aligns with these goals by:

1. **Ensuring Quality Education**: Promoting equitable access to agricultural education for marginalized communities.
2. **Sustainability**: Encouraging research on climate-resilient crops, sustainable farming practices, and resource management.
3. **Global Competitiveness**: Preparing students for global challenges by aligning curricula with international standards.

Social Equity and Agricultural Education

Social equity is a core principle of NEP. It ensures inclusivity by:

1. **Access to Education**: Establishing institutions in rural and underserved areas to bridge educational gaps.
2. **Diversity and Inclusion**: Promoting opportunities for women, marginalized communities, and smallholder farmers in agricultural education.
3. **Financial Support**: Providing scholarships and financial aid to economically disadvantaged students.

Way Forward

To fully realize the potential of NEP in agricultural education, the following steps are essential:

1. **Policy Implementation**: Ensuring effective execution of NEP reforms at institutional and grassroots levels.
2. **Capacity Building**: Training educators and administrators in NEP's multidisciplinary and digital pedagogies.
3. **Public-Private Partnerships**: Encouraging collaboration between educational institutions,

industry, and government to enhance research and innovation.

4. **Global Collaborations**: Establishing partnerships with international universities to share knowledge and best practices.

Conclusion

The New Education Policy provides a transformative vision for agricultural education, addressing contemporary challenges while fostering sustainable and inclusive growth. By aligning educational frameworks with technological advancements and societal needs, the NEP ensures a resilient agricultural sector capable of achieving SDGs and contributing to national development.

References

1. Government of India. (2020). National Education Policy 2020. Ministry of Human Resource Development.
2. Food and Agriculture Organization (FAO). (2015). Sustainable Development Goals and Agriculture.
3. Indian Council of Agricultural Research (ICAR). Annual Reports and Publications.
4. UNESCO. (2020). Education for Sustainable Development: A Roadmap.
5. World Bank. (2021). Transforming Agricultural Education: Global Insights.

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Microorganisms and Minerals Interactions

ARTICLE ID: 0178

B. Himaja[1], M. Manjunath[2], S. Triveni[1], S. Bandeppa[3]

[1]Professor Jayashankar Telangana State Agricultural University, Hyderabad-500030

[2]ICAR-Central Research Institute for Dryland Agriculture, Hyderabad-500059

[3]ICAR-Central Indian Institute of Rice Research, Hyderabad-500030

A mineral is a naturally occurring homogeneous inorganic substance generally in crystalline form with a definite chemical composition. They are usually in solid form except fossil fuels, natural water and mercury (Haldar 2018). Microorganisms interact with minerals which are present in soil. The microorganisms and minerals have coevolved over the years (Li et al 2013). Mineral microbe interactions are important for the supply of nutrients to plant growth and development, biogeochemical cycling of elements, improving quality of environment and formation of ore deposits (Dong et al. 2022).

Role of Microorganisms During Microbe-Mineral Interaction

Mineral weathering

Acidolysis

Various acids such as carbonic acid, gluconic acid, nitric and nitrous acid etc are released by the microorganisms through their metabolic activities to cause dissolution of minerals (Uroz et al. 2009, Dong et al. 2022).

Chelation

Chelating molecules like siderophores and catechol derivatives chelate iron ions and help in dissolution of iron containing minerals such as olivine, hornblende etc. (Uroz et al. 2009; Page et al. 1984; Liermann et al. 2000).

Mineral precipitation

Formation of mineral deposits are significantly influenced by the microorganisms (Hoffmann et al. 2021; Crosby and Bailey 2012). Bacteria help in mineral precipitation by changing local ion concentration, pH or acting as nucleation places for crystallization (Hoffman et al. 2021). Several bacteria, fungi and algae induce carbonate precipitation by

different metabolic activities like ammonification (Lange-Enyedi et al. 2024), sulfate reduction (Zhang 2020), and ureolysis (Hammes et al., 2003).

Mineral Transformation

Microorganisms play important role in formation, dissolution and deterioration of various kinds of minerals (Gadd, 2013). The bacteria such as *Acidithiobacillus ferroxidans*, and *Acidithiobacillus thioxidans* are involved in biooxidation of arsenic (Hackl and Jones 1997). Iron transformation is mediated by *Thiobacillus ferrooxidans*, *Ferrobacillus ferrooxidans*, *Gallionella*, *Leptothrix ochracea*. The sulphur transformation is carried out by *Beggiatoa*, *Thiobacillus*, and *Rhodobacteriineae*. The *Proteus vulgaris* and *Escherichia coli* facilitate transformation of copper compounds (Mulder and Van veen 1968)

Minerals Roles In Helping Microbial Communities

Protection

Minerals help in survival of microbial communities by protecting them from temperatures extremes, uv radiations and physical abrasions through pores, cracks and crevices within minerals (Fomina and Skorochod 2020, Dong et al. 2022).

Nutrients

Minerals provide several nutrients such as phosphorus, potassium, calcium, magnesium, sulfur, iron, sodium etc. for the growth and development of microorganisms (Shock 2009; Li et al 2019; Dong et al. 2022).

Energy

Through oxidation-reduction reactions minerals also act as source of energy to microorganisms (Ehrlich 1996; Shock 2009; Dong et al. 2022).

Uses Of Microbe Mineral Interactions

Mineral solubilizing microorganisms improve plant growth and development by making available nutrients such as P, K and Zn (Devi et al. 2022). Through the process of bioleaching microbe mineral interactions help in extraction of metals like gold and copper from low grade ores and minerals (Bosecker 1997). Application of microbe–mineral suspensions to contaminated areas leads to stabilization of toxic metals either through precipitation or incorporation of metals into biominerals (Dong et al. 2022).

Acknowledgements

The authors thank the authorities of ICAR-CRIDA and PJTSAU, Hyderabad for providing necessary facilities.

References

1. 132 (1–4):5-9, ISSN 0009-2541, https://doi.org/10.1016/S0009-2541(96)00035-6.
2. Bosecker K, 1997. Bioleaching: metal solubilization by microorganisms, *FEMS Microbiology Reviews*, 20 (3-4) :591–604
3. Crosby C.H., Bailey J. 2012 The role of microbes in the formation of modern and ancient phosphatic mineral deposits. Frontiers in Microbiology 3
4. Dong H, Huang L, Zhao L, Zeng Q, Liu X, Sheng Y, Shi L, Wu G, Jiang H, Li F, Zhang L, Guo D, Li G, Hou W, Chen H 2022. A critical review of mineral–microbe interaction and co-evolution: mechanisms and applications, *National Science Review*, 9 (10): nwac128,

5. Ehrlich HL, 1996 How microbes influence mineral growth and dissolution. Chemical Geology,
6. Fomina M, Skorochod I. Microbial Interaction with Clay Minerals and Its Environmental and Biotechnological Implications. *Minerals*. 2020; 10(10):861.
7. Gadd, G.M. (2013). Microbial Roles in Mineral Transformations and Metal Cycling in the Earth's Critical Zone. In: Xu, J., Sparks, D. (eds) Molecular Environmental Soil Science. Progress in Soil Science. Springer, Dordrecht. https://doi.org/10.1007/978-94-007-4177-5_6
8. Haldar S. K.2018. Mineral Exploration, Editor(s): Haldar S.K., Mineral Exploration (Second Edition), Elsevier,2018, Pages 1-23, ISBN 9780128140222, https://doi.org/10.1016/B978-0-12-814022-2.00001-0.
9. Hammes F, Boon N, de Villiers J, Verstraete W, Siciliano SD. Strain-specific ureolytic microbial calcium carbonate precipitation. Appl Environ Microbiol. 2003 Aug;69(8):4901-9. doi: 10.1128/AEM.69.8.4901-4909.2003. PMID: 12902285; PMCID: PMC169139
10. Hoffmann TD, Reeksting BJ, Gebhard S. Bacteria-induced mineral precipitation: a mechanistic review. Microbiology (Reading). 2021 Apr;167(4):001049. doi: 10.1099/mic.0.001049. PMID: 33881981; PMCID: PMC8289221.
11. ISSN 0169-1317
12. Lange-Enyedi, N.T., Németh, P., Borsodi, A.K. et al. Calcium carbonate precipitating extremophilic bacteria in an Alpine ice cave. Sci Rep 14, 2710 (2024). https://doi.org/10.1038/s41598-024-53131-y
13. Li G, Zhou C H, Fiore S, Yu W H 2019. Interactions between microorganisms and clay minerals: New insights and broader applications, Applied Clay Science,177: 91-113
14. Li Jinhua, Benzerara Karim, Bernard Sylvain, Beyssac Olivier 2013. The link between biomineralization and fossilization of bacteria: Insights from field and experimental studies, Chemical Geology, Volume 359, 2013, Pages 49-69, ISSN 0009-2541,
15. Liermann, L.J. et al. (2000) Role of bacterial siderophores in dissolution of hornblende. Geochim. Cosmochim. Acta 64, 587–602
16. Mulder, E. G. and Van veen, W. L. 1968. Effect of microorganisms on the transformation of mineral fractions in soil. Trans. Int. Congr. Soil Sci, 9th 4: 651–661.
17. Page, W.J. and Huyer, M. (1984) Derepression of the *Azotobacter vinelandii* siderophore system, using iron-containing minerals to limit iron repletion. J. Bacteriol. 158, 496–502 52
18. Uroz S, Kelly LC and Turpault M-P et al. The mineralosphere concept: mineralogical control of the distribution and function of mineral-associated bacterial communities. Trends Microbiol 2015; 23: 751–62.
19. Zhang S. 2020. The relationship between organoclastic sulfate reduction and carbonate precipitation/dissolution in marine sediments, Marine Geology, 428:106284, ISSN 0025-3227

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Citrus Decline In Nagpur Mandarin (*Citrus reticulata* Blanco) orchard in South-Eastern Rajasthan

ARTICLE ID: 0179

Himanshu Chawla

Ph.D. Scholar, Department of Fruit Science, College of Horticulture and Forestry, Jhalawar, Agriculture University Kota, Rajasthan – 326023

Mandarins are commercially cultivated in distinct regions across India, including Nagpur Mandarin in Central India, Khasi Mandarin in the Northeast, and Coorg Mandarin in the Southern states. The largest cultivation areas are found in Madhya Pradesh, followed by Maharashtra and Punjab. Among citrus fruits, mandarins account for 42% of the total cultivated area and 43% of the total citrus production. In India, the total mandarin production stands at 6.27 million tons, grown over 0.46 million hectares, yielding a productivity rate of 13.6 tons per hectare (Anon., 2022).

The "Nagpur Mandarin" is considered one of the finest and most popular varieties both in India and globally, renowned for its high-quality fruit. This variety is highly polyembryonic, characterized by medium-sized, upright trees with an evergreen growth habit and relatively few thorns. The leaves are medium-sized, lanceolate in shape, with a prominent midrib and long, narrowly winged petioles. The flowers are small to medium in size, and the fruit is typically globose or sub-globose, with a pale orange-yellow peel that easily separates from the segments. The seeds are small, pointed, and contain green cotyledons, numbering between 10 and 15. Each fruit typically contains 10 to 12 segments. The fruit offers a mild flavour, excellent quality and is notably juicy, with a total soluble solids (TSS) content of 10–12° Brix and an acidity level of 0.5 to 0.9 per cent. Mandarin juice is not only refreshing but also nutritious, appreciated for its ascorbic acid content, sweet-tart taste and vibrant colour.

In Rajasthan, mandarin is the dominant fruit crop in terms of both cultivated area and production. The crop covers an area of 0.24 lakh hectares, yielding

a total production of 0.44 million tonnes (Anon., 2022). Within the state, Jhalawar district leads in mandarin cultivation, particularly of Nagpur mandarin, with cultivation spread over 0.23 lakh hectares and a production of 4.30 lakh tonnes (Anon., 2022). In Jhalawar, growers typically adopt both the Ambe and Mrig bahar cropping patterns, although the Mrig bahar is predominantly favoured. This preference is due to the adverse effects of the high temperatures between April and June, which cause significant fruitlet drop and economic losses for growers. The flowering period for Ambe bahar in Jhalawar lasts from the last week of February to the end of March, with fruit setting completed by the end of March or early April. In contrast, the Mrig bahar flowering season begins in June and continues through the first week of July, with fruit setting completed by the end of July or early August.

Water scarcity significantly impacts mandarin cultivation, with groundwater resources steadily depleting each year. Jhalawar, which receives an annual rainfall of 1200 mm, is also classified as a "dark zone" where water availability is critically low. Water is a crucial component in plants, constituting nearly 90 per cent of their biomass. It comprises 60 to 90 per cent of the root mass, 35 to 95 per cent of the leaves, and 70 to 90 per cent of the water content in fleshy fruits. Consequently, water plays a central role in cytoplasmic functions across all plant tissues. It is also vital for the plant's hydraulic processes and serves as a medium for transporting nutrients from the soil to the plant tissues. During photosynthesis, water is an essential component in the synthesis of food. Additionally, water regulates temperature in plants, acting as a key determinant in their growth, development, and overall survival.

Globally, water scarcity is a pervasive issue affecting all plant-growing regions, with horticulture in Rajasthan and other parts of India being particularly vulnerable. During peak summer, water availability for crop production becomes critically scarce. Over-extraction of groundwater from aquifers, without adequate replenishment, has led to progressively deeper wells. This exacerbates the water shortage, especially in the summer months, leaving no water for both horticulture and agriculture (Mellado *et al*. 2005; Hasan *et al*. 2007).

Citrus decline represents a significant threat in this region, with many orchards experiencing the detrimental effects of this disease. Initially, affected trees exhibit robust growth and perform well during the early years of orchard establishment. However, over time, they begin to deteriorate in health, producing minimal fruit and becoming economically unviable (Koli, 2010). Rather than dying abruptly, the trees undergo a gradual decline, slowly losing vitality. These trees, once a potential source of profit, turn into a considerable liability for growers. The quantity and quality of fruit also suffer markedly, with fruit size shrinking and its overall quality deteriorating. The fruit develops a rough, leathery rind, lacks shine and contains significantly less juice. The feeder root system may become depleted, with roots turning black. In some cases, affected trees persist in a weakened state, slowly declining over time, a condition known as slow decline. However, there are instances where trees die

rapidly within a few days, referred to as quick decline. The disease may affect only a few trees within an orchard or impact the entire orchard within a given locality. The severity of the problem intensifies as the trees age, with trees older than 15 to 20 years becoming increasingly uneconomical. No orchard in the area has been found entirely free from citrus decline (Singh and Bhatnagar, 2017).

References

1. Anonymous. Database, *Agriculture Statistics at a Glance* (2022). Ministry of Agriculture and Farmers Welfare, Govt. of India. 2022;92.
2. Anonymous. Department of Horticulture Report, *Pant Krishi Bhawan,* Government of Rajasthan, Jaipur, 2022.
3. Hasan M, Sirohi NPS, Kumar V, Sharma MK and Singh AK. Performance evaluation of different irrigation scheduling methods for peach through efficient fertigation system network. Acta Horticulturae. 2007;662:193-97.
4. Koli RK. Citrus decline and its management in Rajasthan. Indian Journal of Horticulture. 2010;67(2):203-210.
5. Mellado-Vazquez A, Volke-Haller V, Tapia-Vargas M, Sanchez-Garcia P and Quevedo-Nolasco A. Response of papaya to irrigation and N-P-K fertilization in a vertisol. Terra. 2005;23:137-44.
6. Singh B, and Bhatnagar P. Impact of climate change on citrus decline in India. Journal of Agricultural Research and Development. 2017;48(1):50-56.

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Fungi for the Future: Revolutionizing Agriculture with Mushroom Cultivation

ARTICLE ID: 0180

Juhi Ranjan[1]*, Dr. Satvaan Singh[2]

[1]Student, Department of Agriculture, Keral Verma Subharti College of Science, Swami Vivekanand Subharti University, Meerut-250005, Uttar Pradesh, India

[2]Assistant Professor, Department of Agriculture, Keral Verma Subharti College of Science, Swami Vivekanand Subharti University, Meerut-250005, Uttar Pradesh, India

Mushroom cultivation is a fast-growing agricultural sector with immense potential for addressing food security, promoting sustainability, and generating income. As a nutritious food source and a valuable ingredient in medicine and industry, mushrooms are gaining popularity worldwide.

The Importance of Mushroom Cultivation

Mushrooms are a rich source of protein, vitamins, minerals, and antioxidants. They contain essential nutrients such as B vitamins, selenium, and potassium while being low in calories and fat. Their unique umami flavour makes them a versatile ingredient in various cuisines. Beyond nutrition, mushrooms are used in traditional and modern medicine for their immune-boosting, anti-inflammatory, and anti-cancer properties.

Mushroom cultivation is also an environmentally friendly practice. Unlike conventional crops, mushrooms require minimal land, water, and energy to grow. They thrive on agricultural and industrial waste, converting it into valuable food products. This makes mushroom farming a key player in circular economies and waste management strategies.

The Basics of Mushroom Cultivation

Successful mushroom cultivation depends on understanding the lifecycle of fungi and creating optimal growing conditions. The process involves several key steps:

1. Selecting the Mushroom Variety

The choice of mushroom species depends on local climate, available resources, and market demand. Commonly cultivated mushrooms include:

Button Mushrooms (*Agaricus bisporus*): Popular and easy to grow.

Oyster Mushrooms (*Pleurotus spp.*): Fast-growing and adaptable to various substrates.

Shiitake Mushrooms (*Lentinula edodes*): Valued for their medicinal properties.

Enoki Mushrooms (*Flammulina velutipes*): Grown for their delicate flavor and texture.

2. Preparing the Substrate

The substrate serves as the growth medium for mushrooms. It typically consists of organic materials such as straw, sawdust, or agricultural by-products. The substrate must be sterilized or pasteurized to eliminate competing microorganisms.

3. Inoculation

Inoculation involves introducing mushroom spawn (mycelium) into the substrate. Spawn acts as the "seed" for mushroom cultivation and is typically produced in specialized laboratories to ensure quality and purity.

4. Incubation

During this phase, the inoculated substrate is kept in a controlled environment with optimal temperature, humidity, and darkness. The mycelium colonizes the substrate, forming a network of fungal threads.

5. Fruiting

Once the substrate is fully colonized, environmental conditions are adjusted to trigger fruiting. This includes introducing fresh air, light, and maintaining high humidity. Mushrooms begin to emerge within a few days.

6. Harvesting and Post-Harvest Handling

Mushrooms are harvested at their peak maturity to ensure freshness and quality. Proper handling, cooling, and packaging are crucial to maintaining their shelf life and market value.

Methods of Mushroom Cultivation

Several cultivation methods cater to different scales and resource availability. These include:

1. Indoor Farming

Indoor cultivation involves growing mushrooms in controlled environments such as grow rooms, greenhouses, or containers. This method allows year-round production and precise control over growing conditions. It is ideal for commercial operations targeting consistent quality and quantity.

2. Outdoor Farming

Outdoor cultivation mimics natural growing conditions and is typically used for species like shiitake and reishi. Logs, stumps, or shaded beds serve as substrates, and minimal infrastructure is required. However, this method is seasonal and dependent on climate.

3. Bag Cultivation

Bag cultivation is a versatile technique where substrates are packed into plastic bags and inoculated with spawn. The bags act as mini grow chambers, providing the necessary environment for mycelium growth and fruiting.

4. Hydroponic Systems

Advanced hydroponic systems use nutrient-rich water solutions and soilless substrates to grow mushrooms. While less common, this method offers potential for urban farming and innovative designs.

Benefits of Mushroom Cultivation

1. Economic Opportunities

Mushroom farming provides lucrative opportunities for small-scale and commercial growers. With high market demand and relatively low production costs, mushrooms offer attractive profit margins. Value-added products such as dried mushrooms, mushroom powders, and processed foods further enhance income potential.

2. Sustainability

By utilizing agricultural waste as substrates, mushroom cultivation promotes recycling and reduces landfill waste. It also has a smaller carbon footprint compared to conventional agriculture.

3. Food Security

Mushrooms are a quick and reliable food source that can be grown in diverse environments. They contribute to dietary diversity and address malnutrition in resource-limited regions.

4. Environmental Benefits

Mushrooms play a role in bioremediation by breaking down pollutants and improving soil health. Their ability to degrade complex organic compounds makes them valuable in waste treatment projects.

Challenges in Mushroom Cultivation

Despite its advantages, mushroom farming faces several challenges:

1. Technical Expertise

Successful cultivation requires knowledge of fungal biology, environmental controls, and pest management. Training and resources are essential for new growers.

2. Market Access

Limited awareness and market infrastructure can hinder the sale of mushrooms, particularly in rural areas. Building distribution networks and educating consumers are key to expanding market reach.

3. Infrastructure and Investment

Establishing controlled environments and acquiring quality spawn can be expensive for small-scale farmers. Financial support and subsidies can alleviate these barriers.

4. Pest and Disease Management

Fungi are susceptible to contamination by competing molds, bacteria, and pests. Maintaining hygiene and monitoring growing conditions are crucial to prevent losses.

Global Trends in Mushroom Cultivation

The mushroom industry is witnessing significant growth and innovation worldwide. Key trends include:

1. Organic and Sustainable Farming

Consumers are increasingly demanding organic mushrooms grown without synthetic chemicals. Sustainable practices such as biodegradable packaging and renewable energy are gaining traction.

2. Medicinal Mushrooms

Interest in medicinal mushrooms like reishi, lion's mane, and cordyceps is rising due to their health benefits. These mushrooms are being integrated into supplements, teas, and skincare products.

3. Urban Farming

Urban mushroom farms are emerging in cities, utilizing unused spaces like basements and rooftops. These farms cater to local markets and reduce transportation emissions.

4. Technological Innovations

Automation, AI-driven climate control systems, and advanced spawn production techniques are revolutionizing mushroom farming. These innovations improve efficiency and scalability.

Conclusion

Mushroom cultivation is a promising and sustainable agricultural practice that offers nutritional, economic, and environmental benefits. By understanding the science and methods of cultivation, individuals and communities can tap into the potential of fungi farming. Addressing challenges through education, infrastructure, and innovation will further unlock the industry's growth.

As global interest in mushrooms continues to rise, fostering a culture of sustainable mushroom farming will contribute to food security, environmental health, and economic prosperity. Whether as a backyard hobby or a commercial venture, mushroom cultivation is an invaluable tool for creating a better future.

References

1. Chang, S.T., and Miles, P.G. "Mushrooms: Cultivation, Nutritional Value, Medicinal Effect, and Environmental Impact." CRC Press, 2004.
2. Singh, M., Vijay, B., Kamal, S., and Wakchaure, G.C. "Mushroom Cultivation, Marketing and Consumption." Directorate of Mushroom Research, Solan, India.
3. United Nations Food and Agriculture Organization (FAO). "Mushrooms and their Role in Sustainable Development."

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Battling the Bugs: A Comprehensive Guide to Managing Insect Pests in Mustard Cultivation

ARTICLE ID: 0181

Sahil Shamkuwar[1*], Rutuja Tayade[2]

[1]Department of Genetics and Plant Breeding, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi

[2]Department of Agricultural Entomology, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi

Mustard (*Brassica spp.*) is an essential global oilseed crop known for its adaptability and resilience in diverse climatic conditions. Primarily cultivated for its seeds, which contain 30-45% oil and 20-25% protein, mustard significantly contributes to agricultural practices by enhancing soil fertility and reducing pests and diseases through its allelopathic properties (Bahl *et al.,* 2021). It is considered sustainable, thriving in suboptimal soils with minimal inputs, thus supporting food security and agricultural diversity (Mishra *et al.,* 2020). Mustard cultivation supports millions of farmers, particularly in India, Canada, and the EU, serving as a vital income source.

Despite of global importance, mustard possess serious threat due various biotic and abiotic factors. There are various factors which limits the mustard production in the country and needs the critical attention towards them. These factors are non-adoption of improved package of practices and production technologies, susceptibility of mustard varieties to various pest and diseases and unmanaged crop growth. As every crop has to suffer from biotic and abiotic stresses, mustard is likely to get affected by insect pest like Aphids, Painted bug, Diamondback moth, Mustard Sawfly which are considered to be a major pest in mustard and minor pest includes, Green Peach Aphid, Pea Leaf-minor, Bihar Hairy Caterpillar and Cabbage butterfly. These pests threaten the crop right from sowing until end of crop season. These insect-pest can cause yield losses of 20% to 50% by damaging plants (Singh *et al.,* 2022). Additionally, pests can compromise seed quality by causing discoloration and

reducing oil content while increasing susceptibility to diseases.

Need of the Hour

Current research trends focus on breeding high-yielding and disease-resistant varieties through genomics (Kumar *et al.,* 2023) and developing integrated pest management (IPM) practices to reduce reliance on chemical pesticides while improving yield and quality (Singh *et al.,* 2022). To address these challenges, IPM strategies are essential for effectively managing pest populations while minimizing environmental impact. These strategies may include cultural practices such as crop rotation and the use of natural predators. Overall, effective pest management is crucial for maximizing mustard yield and quality while promoting sustainable agricultural practices that protect both the environment and public health.

Insect-pest of Mustard

Mustard crops face several significant insect pests that can adversely affect their yield and quality. The major pests include:

1. Mustard Aphid

a. **Scientific name:** *Lipaphis erysimi,* **Family:** Aphididae, **Order:** Hemiptera

b. **Life stages and identification:** Aphids are small, soft-bodied, pearl-shaped insects that have a pair of cornicles.

c. **Symptoms and nature of Damage:** Both nymphs and adults suck sap from leaves, buds, and pods, causing yellowing, curling, and drying of leaves. This leads to weak pods and small seeds. They also secrete honeydew, which promotes sooty mold and reduces photosynthesis.

d. **Losses caused to plant:** When the ETL is crossed, the losses can range from 20-45% under natural field condition.

2. Mustard Saw Fly

a. **Scientific name:** *Athalia lugens proxima,* **Family:** Tethredinidae, **Order:** Hymenoptera

b. **Life stages and identification:** The larva is greenish black and feeds on mustard leaves during the morning and evening and make holes. The adult insect is small, orange-yellow, and features black markings and smoky black veins.

c. **Symptoms and nature of Damage:** Grubs alone are destructive, bite holes into leaves preferring young growth and skeletonize the leaves completely. When infested mature plants, seed setting is affected.

d. **Losses caused to plant:** Mustard sawfly can cause damage from 5 to 18 % as reported in many parts of the country.

3. Painted Bug

a. **Scientific name:** *Bagrada hilaris,* **Family**: Pentatomidae, **Order:** Hemiptera
b. **Life stages and identification:** The adult is black with orange spots, red and yellow markings when turn to adult.
c. **Symptoms and nature of Damage:** Nymphs and adult bugs suck sap from leaves and developing pods, causing young plants to wilt and dry up, while also excreting a resinous material that spoils the pods; infestation at maturity leads to pod curling and leaf shriveling.
d. **Losses caused to plant:** Painted bugs reduce the mustard crop yield by 30-40%, causing up to 50% pod damage and decreases seed quality.

4. **Diamondback Moth**

a. **Scientific name:** *Plutella xylostella,* **Family**: Plutellidae, **Order:** Lepidoptera
b. **Life stages and identification:** Larva is yellowish green, with fine erect black hairs scattered all over the body and adult are small greyish brown having pale whitish narrow wings with yellow inner margins.
c. **Symptoms and nature of Damage:** Larvae cause whitish patches on leaves by scraping epidermal tissues, giving them a withered appearance, and later bore holes. They can completely consume leaves and also damage pods by feeding on developing seeds.
d. **Losses caused to plant:** Diamondback moths can cause up to 40-50% yield loss in mustard, with severe infestations damaging up to 60% of leaves and affecting pod quality.

Integrated Pest Management (IPM) Strategies

IPM is a systems approach that combines a wide array of crop production and protection measures to minimize the economic losses caused by pest. Hence, use of low or judicious dose of pesticides, integrated with other means like growing pest tolerant cultivars, sanitation, crop rotation, use of bio-agents and plant extracts seems to be best method of pest management without environmental pollution.

Insect-pest	Management
Aphids	• Early sowing, preferably up to third week of October. • Apply any of the following insecticides when pest levels reach 50-60 aphids per 10 cm of the central shoot, or when 40-50% of the plants are infested. For foliar sprays, use 625-1000 ml of oxydemeton-methyl 25 EC, dimethoate 30 EC, , quinalphos 25 EC, or malathion 50 EC; or 940-1500 ml of chlorpyrifos 20 EC in 600-1000 L of water per hectare, depending on the crop stage. For granular

	insecticides, apply 10 kg of phorate 10 G or 33 kg of carbofuran 30 per hectare, followed by light irrigation.
Mustard Saw Fly	• Early sowing • Administer the first irrigation 3-4 weeks after sowing to significantly lower the bug population. • Apply 1.0 L of malathion 50 EC or quinalphos 25 EC mixed in 500-600 L of water per hectare once in October and again in March-April.
Painted Bug	• Early sowing. • Spray the crop with malathion 50 EC @1000 ml or dimethoate 30EC @ 625 ml in 600- 700 liter water
Diamondback Moth	• For control of grown up larvae apply 5% malathion dust @ 37.5 kg/ha

To effectively manage insect pests in mustard crops, a variety of strategies can be applied. Start by selecting tolerant varieties like JM-1 and RK-9501 for aphids, and aim to sow the crops before October 20 to reduce damage. Use yellow stick traps to monitor aphid populations and promptly remove affected plant parts. It's important to conserve beneficial insects such as ladybird beetles (*Coccinella septempunctata*, *Menochilus sexmaculata*, *Hippodamia variegata*, and *Cheilomones vicina*), which can consume 10 to 15 aphids each day, along with syrphid flies (*Sphaerophoria spp.*, *Eristallis spp.*, *Metasyrphis spp.*, *Xanthogramma spp*., and *Syrphus spp.*). The braconid parasitoid *Diaretiella rapae* and lacewing (*Chrysoperla carnea*) also play a significant role in controlling aphid populations.

For managing mustard sawflies, practice summer ploughing to eliminate pupae, maintain clean cultivation practices, and ensure proper irrigation during the seedling stage to drown larvae. Regularly collect and destroy sawfly grubs. To address painted bugs, deep ploughing can help eliminate their eggs, while early sowing and irrigation during the first four weeks after planting can mitigate pest pressure. Quick threshing of harvested crops and burning leftover crop residues will prevent pests from surviving into the next season.

For diamondback moths, install pheromone traps to keep track of their activity and regularly collect larvae. It is also beneficial to conserve parasitoids like *Cotesia plutellae* and *Diadegma insulare*, and apply malathion dust for controlling larger larvae. Additionally, using bitter gourd seed oil emulsion can serve as an anti-feedant. For chemical control options, consider spraying the crop with malathion 50 EC or quinolphos at the recommended concentrations mixed with water. By implementing these integrated pest management techniques, you can significantly reduce pest populations and safeguard mustard crops.

Conclusion

In conclusion, effective insect pest management in mustard cultivation is crucial for enhancing crop efficiency and production. The integration of various strategies within an Integrated Pest Management (IPM) framework allows for a sustainable approach to controlling pest populations while minimizing environmental impact. Recently developed technologies insures management of insect-pest very reliable with reduced cost and environment friendly.

References

1. Bahl, P., Kaur, S., & Kumar, S. (2021). Nutritional evaluation of mustard (Brassica spp.) seeds: A review. Journal of Agricultural and Food Chemistry, 69(3), 765-775.
2. Kumar, V., Yadav, A., & Singh, P. (2023). Advances in mustard breeding: Genomics and biotechnology approaches. Field Crops Research, 292, 108764.
3. Mishra, A., Rani, S., & Singh, S. (2020). Sustainable agriculture: Role of mustard in crop rotation. Agricultural Reviews, 41(2), 152-160.
4. Singh, R., Gupta, A., & Singh, D. (2022). Integrated pest management in mustard: Current trends and future prospects. Journal of Integrated Pest Management, 13(1), 1-10

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Nursery Bed Preparation for Flower Cultivation

ARTICLE ID: 0182

Dr. Satvaan Singh[1]*, Juhi Ranjan[2]

[1]Assistant Professor, Department of Agriculture, Keral Verma Subharti College of Science,Swami Vivekanand Subharti University, Meerut-250005, Uttar Pradesh, India

[2]Student, Department of Agriculture, Keral Verma Subharti College of Science,Swami Vivekanand Subharti University, Meerut-250005, Uttar Pradesh, India

Nursery bed preparation is a fundamental step in flower cultivation, ensuring the healthy growth of seedlings and the success of flower farming. A well-prepared nursery bed provides optimal conditions for germination, root development, and protection against pests and diseases.

The Importance of Nursery Bed Preparation

A nursery bed serves as the starting point for cultivating strong and healthy seedlings. Proper preparation of nursery beds ensures:

1. **Healthy Germination:** Providing a conducive environment for seeds to sprout and grow.
2. **Pest and Disease Management:** Reducing the risk of infections and pest infestations by creating sterile conditions.
3. **Optimal Root Development:** Offering loose, nutrient-rich soil that promotes robust root systems.
4. **Efficient Resource Use:** Conserving water, nutrients, and space by concentrating resources on young plants.
5. **Easy Transplantation:** Ensuring seedlings are strong enough to survive transplantation with minimal shock.

Types of Nursery Beds

There are three primary types of nursery beds, each suited to specific needs and environmental conditions:

1. **Raised Beds:** Elevated beds that improve drainage and are ideal for regions with heavy rainfall.
2. **Sunken Beds:** Beds that are slightly below ground level, suitable for dry areas as they retain moisture.

3. **Flat Beds:** Level beds used in areas with moderate climatic conditions, suitable for most flower varieties.

The choice of nursery bed type depends on local climate, soil conditions, and the specific requirements of the flowers being grown.

Steps in Nursery Bed Preparation

The process of preparing a nursery bed involves several crucial steps:

1. **Site Selection**

- Choose a location with adequate sunlight, good drainage, and protection from strong winds.
- Ensure proximity to a water source for easy irrigation.

2. **Soil Preparation**

- **Soil Testing:** Conduct soil tests to determine pH, nutrient levels, and texture. Most flowers thrive in slightly acidic to neutral soil (pH 6.0–7.5).
- **Loosening the Soil:** Dig the soil to a depth of 20–30 cm to break up compacted layers and improve aeration.
- **Adding Organic Matter:** Incorporate well-decomposed compost, farmyard manure, or leaf mold to enrich the soil with nutrients.
- **Sterilization:** Sterilize the soil to eliminate pathogens and weed seeds. This can be done through solarization (covering the bed with plastic sheets under sunlight) or by using chemical sterilizers.

3. **Bed Shaping**

- Level the soil and shape the bed according to the chosen type (raised, sunken, or flat).
- Maintain a width of 1–1.5 meters for easy access, with pathways in between for movement.
- Ensure the bed has a gentle slope for proper drainage.

4. **Seed Sowing**

- Create furrows or drills at appropriate depths for the specific flower seeds.
- Sow seeds at the recommended spacing to prevent overcrowding.
- Cover seeds lightly with a thin layer of soil or sand to protect them from birds and pests.

5. **Watering**

- Water the bed gently using a fine sprinkler to avoid displacing seeds.
- Maintain consistent moisture levels during the germination period.

6. **Mulching**

Apply a thin layer of organic mulch (e.g., straw or grass clippings) to retain soil moisture, regulate temperature, and suppress weeds.

Special Considerations for Flower Nursery Beds

1. **Flower-Specific Needs**

Different flower species have unique requirements for soil, light, and moisture. For instance:

Marigolds: Prefer well-drained loamy soil and full sunlight.

Roses: Thrive in slightly acidic soil enriched with organic matter.

Petunias: Require light, sandy soil with good drainage.

Understanding the specific needs of the flowers being cultivated is crucial for successful nursery bed preparation.

2. **Integrated Pest Management (IPM)**
   - Use biological controls such as neem oil or predatory insects to manage pests.
   - Avoid overcrowding to reduce the risk of fungal infections.
3. **Seasonal Timing**
   - Prepare nursery beds a few weeks before sowing to allow soil to settle and nutrients to integrate.
   - Align sowing with the flowering season to ensure blooms at the right time.

Maintenance of Nursery Beds

Proper maintenance of nursery beds ensures the healthy growth of seedlings:

1. **Weeding:** Remove weeds regularly to prevent competition for nutrients and water.
2. **Irrigation:** Maintain consistent moisture levels without waterlogging.
3. **Thinning:** Remove excess seedlings to allow adequate space for growth.
4. **Fertilization:** Apply balanced fertilizers or organic compost during the growth phase to support plant development.
5. **Shading:** Use shade nets or covers to protect young seedlings from extreme weather conditions.

Advantages of Well-Prepared Nursery Beds

1. **Higher Germination Rates:** Properly prepared beds provide optimal conditions for seed germination.
2. **Uniform Growth:** Healthy, evenly spaced seedlings develop into uniform plants.
3. **Reduced Mortality:** Strong seedlings are better equipped to survive transplantation and resist stress.
4. **Efficient Resource Use:** Concentrating resources in a nursery bed reduces wastage and maximizes productivity.

Challenges in Nursery Bed Preparation

1. **Soil Quality Issues:** Poor soil fertility or contamination can hinder seedling growth.
2. **Pest and Disease Outbreaks:** Inadequate sterilization or overcrowding increases susceptibility to pests and diseases.
3. **Climate Variability:** Extreme weather conditions such as droughts or heavy rains can disrupt the nursery environment.
4. **Labor Intensity:** Preparing and maintaining nursery beds requires significant manual effort and attention to detail.

Future Trends in Nursery Bed Preparation

1. **Automation and Mechanization:** Tools and machines for soil preparation, sowing, and irrigation are making nursery bed management more efficient.
2. **Sustainable Practices:** Organic amendments, bio-fertilizers, and renewable mulching materials are gaining popularity.
3. **Smart Farming:** IoT devices and sensors are being used to monitor soil moisture, temperature, and nutrient levels in real time.
4. **Urban Gardening:** Compact nursery bed systems are being designed for small spaces and rooftop gardens.

Conclusion

Nursery bed preparation is the cornerstone of successful flower cultivation, providing a strong foundation for healthy and vibrant blooms. By

understanding the science and techniques of nursery bed preparation, gardeners and farmers can optimize germination, growth, and yield. As agriculture embraces sustainability and technology, nursery bed practices are evolving to meet the demands of modern flower farming. By investing time and effort in preparing nursery beds, growers can ensure a flourishing garden or a profitable commercial venture. The beauty of flowers begins in the nursery bed, where careful preparation sets the stage for nature's most exquisite creations.

References

1. United Nations Food and Agriculture Organization (FAO). "Sustainable Practices in Horticulture."
2. Indian Council of Agricultural Research (ICAR). "Best Practices for Flower Nursery Management."
3. Royal Horticultural Society (RHS). "Guide to Seed Sowing and Propagation."

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Osmoregulation in Freshwater and Marine Water Fishes

ARTICLE ID: 0183

Arvind Sinha[1], Amit Singh Kshatri[2*], Ramjas Choudhary[1], Divyani Soni[3], Gurudev Ghidorey[4], Ambrish Singh[5]

[1]M.F.Sc., Department of Fisheries Resource Management, College of Fisheries Science, CCS HAU, Hisar, Haryana (125004) India

[2*]PhD Scholar, Department of Fisheries Resource Management, College of Fisheries Science, CCS HAU, Hisar, Haryana (125004) India

[3]M.F.Sc., Department of Fisheries Extension, Late Shri Punaram Nishad College of Fisheries Kawardha, DSVCKV, Durg, Chhattisgarh 491995

[4]M.F.Sc., Department of Fisheries Extension, College of Fisheries Science, CCS HAU, Hisar, Haryana (125004) India

[5]Assistant Professor, Department of Fisheries Resource Management, College of Fisheries Science, Uttar Pradesh Pandit Deen Dayal Upadhyay Pasu Chikitsa Vigyan Vishwavidyalaya and Goo Anusandhan Sansthan, Mathura -281001

The osmoregulatory mechanism in the body of fish adjusts the internal environment to the changing water content of the external environment. The alterations of the internal medium are performed by the coordinated functioning of the kidney, gills, and integument. In seawater fish, the gills and the kidneys play an important role in water retention by concentrating urine of relatively high osmolality. However, in freshwater fish, the kidney is more important in the excretion of the water taken from the hypertonic environment. The concentration of urine is low in the relatively hypotonic environment. In freshwater fish, through the functional activity of the kidney, the maintenance of the blood and tissues in a hypoosmotic climate and a good compensatory mechanism through the gills is

necessary—these mechanisms help to maintain acid-base balance in both conditions. Furthermore, the acid-base equilibrium is of greater importance in freshwater than in seawater fish. In freshwater, the increase in the influx of alkali ions occurs. These ions have to be transported in both directions through the gills. There is a loss of H^+ through an increase in CO_2 excretion only (the main system to excrete CO_2 is through the kidney). There are also effects on ammonia excretion, or ammonia is excreted in the undissociated form, i.e., by passive diffusion through the gills. Ammonia also changes from the undissociated form to the ammonium ion by an increase in NH_3.

Background of Osmoregulation In Fishes

The evolutionary history of osmoregulation in fishes traces back to the earliest vertebrates, which are believed to have originated in marine environments during the Cambrian period, over 500 million years ago. Early fishes likely developed osmoregulatory mechanisms to cope with the challenges posed by their surroundings. Over time, as fishes diversified and adapted to different aquatic environments, these mechanisms evolved to suit varying osmotic conditions.

Fishes are broadly categorized into marine and freshwater species, each facing unique osmoregulatory challenges:

1. **Marine Fishes:** These live in an environment with high salinity (hypertonic), where the concentration of salts in the water is greater than in their body fluids. Marine fishes face the risk of dehydration as water tends to move out of their bodies through osmosis.
2. **Freshwater Fishes:** These inhabit a low-salinity (hypotonic) environment where the salt concentration in the water is much lower than in their body fluids. Freshwater fishes are at risk of swelling and losing salts due to water influx through osmosis.
3. **Euryhaline Fishes:** Some species, like salmon, can tolerate a wide range of salinities and migrate between freshwater and seawater. They exhibit remarkable osmoregulatory flexibility.

Importance of Osmoregulation in Fishes

1. **Maintaining Cellular Function**: The balance of water and salts is essential for proper cellular processes, including enzyme activity, nutrient transport, and nerve function.
2. **Survival in Diverse Habitats**: Osmoregulation allows fishes to thrive in a wide variety of aquatic environments, from freshwater rivers and lakes to marine ecosystems and brackish waters.
3. **Adaptations to Extreme Environments**: Some fishes, such as those in hypersaline or ion-deficient waters, exhibit specialized osmoregulatory mechanisms to survive in extreme conditions. These adaptations include salt-excreting glands, specialized gills, and efficient kidneys.
4. **Energy Efficiency**: Effective osmoregulation minimizes energy expenditure by maintaining an optimal internal environment despite external fluctuations.
5. **Reproductive Success**: Proper osmoregulatory function is critical for reproductive processes, particularly for species that spawn in waters with salinity levels different from their usual habitat.

6. **Evolutionary Adaptations**: Osmoregulation has played a key role in the evolutionary success and diversification of fishes, contributing to their ability to colonize a wide range of ecological niches.

Osmoregulation In Freshwater Fishes

The fish gills and oral membrane allow them to continuously absorb water through diffusion because their internal environment is more concentrated than their surroundings (hyperosmotic to the surrounding medium). Large volumes of diluted urine are continuously expelled by well-developed kidneys as excess water. The pituitary hormones (Example: Arginine-Vasotocin; AVT) that cause blood pressure changes have an impact on the regulation of diuretic (urine-producing) processes. The kidneys of freshwater teleost have many glomeruli and are incredibly large for their body size. Additionally, the glomeruli are large enough to allow for the production of a significant amount of glomerular ultrafiltrate and, subsequently, a large volume of urine. Due to the large amount of urine they secrete, many freshwater teleost have developed a urinary. Despite the urine's low salt content, a considerable amount of salt is lost due to its frequent flow (Greenwell *et al.,* 2003). Additionally, salts are lost from the gill tissues through diffusion. Consuming salt in food and actively absorbing it through the gills balance these losses. Sodium and chloride pumps are found in specific cells found in gill lamellae. In order to maintain the ions' higher concentration in the body, these pumps—special enzymes—use energy to move the ions up their concentration gradient.

Osmoregulation In Marine Water Fishes

These fish are at risk of losing water from their bodies because the salt concentrations in their internal environment are about one-third that of their surroundings (they are hypo-osmotic to the surrounding water). It has been observed that osmosis causes some species to lose between 30 and 60% of the water they consume. The skin of marine teleost is impermeable to both water and salts to minimize water loss (Whittamore, 2012). In marine teleost, the quantity and size of glomeruli are either drastically diminished or eliminated. In addition, there isn't a water-impermeable distal convoluted tubule to save water. In addition, the glomerular filtration rate is significantly decreased, and less urine is produced to prevent the loss of water. As a result, freshwater teleost urine is much less concentrated than that of marine teleost.

In the past, marine teleost would consume a lot of seawater to stay hydrated. The rate of drinking varies by species and by salinity within species. The amount of seawater that marine species typically consume each day ranges from 7 to more than 35% of their body weight. This leads to consuming a lot of salt, which raises the body's salt content.



Figure: Osmoregulation in **A)** Freshwater Fish and **B)** Marine Water Fish

Kidney Function And Osmoregulation

Fish have evolved specialized kidney functions to address osmoregulatory challenges in both freshwater and marine environments. In freshwater, fish face the challenge of too much water entering their bodies because their body fluids are saltier than the surrounding water. To maintain osmotic balance, they produce large amounts of dilute urine, effectively removing the excess water absorbed through their skin and gills, this process helps prevent cellular swelling and ensures the fish's internal salt concentrations remain stable.

Marine fish face the challenge of osmoregulation in hypertonic seawater, where water loss can lead to dehydration. To manage this, their kidneys excrete divalent ions like calcium and magnesium, while gill ionocytes actively secrete monovalent ions such as sodium and chloride. Additionally, to conserve water, marine fish produce minimal, concentrated urine, as their kidneys cannot produce urine more concentrated than their body fluids. These renal adaptations are crucial for maintaining homeostasis in varying aquatic environments.

In a variety of aquatic settings, fish display a range of behavioural adaptations to cope with osmoregulatory challenges. For instance, euryhaline species like the short-finned molly (*Poecilia sphenops*) inhabit estuaries with varying salinities. Their capacity to withstand a broad range of salinities enables them to flourish in these fluctuating environments.

Behavioral Adaptation

In order to spawn, some species, such as salmon, migrate from the ocean to freshwater rivers. In order to adapt to the varying salinities they encounter throughout their life cycle, this migration necessitates significant physiological and behavioural changes (Lillywhite & Evans 2021).

Certain fish modify their behaviour to steer clear of harsh environmental circumstances that might interfere with osmoregulation. For example, some species might become more active at night in order to avoid the high temperatures and evaporation rates that cause salinity to rise during the day. These behavioural strategies are crucial for fish to maintain osmotic balance, ensuring their survival and reproductive success in environments with varying salinity levels.

Conclusion

Osmoregulation is a vital process that allows freshwater and marine fish to survive and thrive in environments with drastically different salinity levels. Freshwater fish face the challenge of excessive water entering their bodies and must excrete large amounts of dilute urine to maintain balance, while marine fish must counteract water loss by drinking seawater and excreting excess salts. These adaptations highlight the remarkable physiological mechanisms fish have developed to regulate their internal environment, ensuring survival in their respective habitats. Understanding osmoregulation not only deepens our knowledge of aquatic life but also provides insights into broader ecological and evolutionary processes.

References

1. Greenwell, M. G., Sherrill, J., & Clayton, L. A. (2003). Osmoregulation in fish: mechanisms and clinical implications. *Veterinary Clinics: Exotic Animal Practice*, *6*(1), 169-189.
2. Lillywhite, H. B., & Evans, D. H. (2021). Osmoregulation by vertebrates in aquatic environments. *eLS*, *2*, 1-14.
3. Whittamore, J. M. (2012). Osmoregulation and epithelial water transport: lessons from the intestine of marine teleost fish. *Journal of Comparative Physiology B*, *182*(1), 1-39.

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Genetic Mechanisms of Flowering Synchronization in Carrot Seed Production

ARTICLE ID: 0184

Jyoti Verma[1], Navdeep Singh[1], Dilpreet Talwar[2]

[1]Research Scholar, Department of Vegetable Science, College of Horticulture and Forestry, Punjab Agricultural University, Ludhiana, Punjab, India

[2]Extension Scientist, Department of Vegetable Science, College of Horticulture and Forestry, PAU, Ludhiana, Punjab, India

Synchronizing flowering is essential in carrot (*Daucus carota* L.) seed production, especially for hybrid seed systems that require precise timing between male-sterile and pollinator lines. Achieving uniform flowering is crucial for successful pollination, maximizing seed yield, and ensuring high seed quality. In hybrid seed production, proper synchronization is particularly important to prevent mismatched pollination periods, which can reduce seed set. Furthermore, the demand for high-quality and genetically pure seeds highlights the necessity of controlling flowering time effectively. However, several factors make achieving consistent flowering across different genotypes and environments challenging. One major obstacle to uniform flowering is genetic variability, as different carrot varieties respond differently to flowering cues. Additionally, plant hormones such as gibberellins and abscisic acid influence flowering, adding another layer of complexity. Agronomic factors, including planting dates, plant density, and irrigation practices, also contribute to variations in flowering time within a seed crop.

Genetic Control of Flowering Synchronization in Carrot

Flowering synchronization in carrots is regulated by genetic mechanisms that integrate environmental signals to ensure uniform and timely floral transition. This regulation is particularly important in seed production systems, where male-sterile and pollinator lines must flower simultaneously for efficient cross-

pollination. Several key genes and molecular pathways contribute to flowering control, responding to factors such as vernalization (cold exposure) and photoperiod (day length). Additionally, epigenetic modifications play a role in regulating flowering gene expression, ensuring that the plant transitions to reproductive development under favourable conditions.

Major Genes and Pathways Involved in Flowering Regulation

Flowering in carrots is controlled by a network of genes that interact with environmental signals to determine when floral initiation occurs. Among these, the FLOWERING LOCUS T (*FT*) and LEAFY (*LFY*) genes play fundamental roles in regulating flowering time and floral organ formation.

Key Flowering Genes Identified in Carrots:

1. Flowering Locus T (*DcFT*)

- The *DcFT* gene in carrots is an important regulator of flowering, functioning similarly to the FT gene in other plant species.
- It acts as a florigen, meaning it produces a protein signal that travels from the leaves to the shoot apical meristem, initiating the transition from vegetative to reproductive growth.
- The expression of *DcFT* is influenced by both photoperiod and vernalization, ensuring that flowering occurs under appropriate environmental conditions.
- Experimental studies indicate that overexpression of *DcFT* leads to early flowering, confirming its critical role in floral induction.

2. Leafy (*DcLFY*)

- The *DcLFY* gene is responsible for defining floral meristem identity, ensuring that flower structures develop properly.
- Once DcFT triggers floral initiation, *DcLFY* helps transition the meristem into reproductive development.
- This gene ensures the correct formation of floral organs, making it essential for proper flowering and reproductive success.

3. Other Flowering-Related Genes in Carrots

- DcSOC1 (SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1): Acts as a central integrator of vernalization and photoperiod signals, helping regulate *DcFT* expression.
- *DcFLC* (FLOWERING LOCUS C-like genes): Functions as a floral repressor, preventing premature flowering before vernalization occurs. Cold exposure reduces *DcFLC* activity, allowing flowering to proceed.
- *DcGI* (GIGANTEA): Plays a role in the circadian clock system and influences *DcFT* expression under long-day conditions.

Comparison of Carrot Flowering Genes with Model Plants

Many flowering genes in carrots have homologs in well-researched model plants, such as Arabidopsis thaliana. By comparing carrot flowering genes with those in Arabidopsis, researchers have gained insights into their functions and interactions.

While carrots share several flowering-related genes with Arabidopsis, differences exist in their responses to environmental cues. Carrots exhibit a stronger dependence on vernalization for flowering,

highlighting the need for further research into their unique genetic regulation mechanisms.

Gene	Function in Arabidopsis	Function in Carrot
FT (FLOWERING LOCUS T)	Produces florigen to initiate flowering	DcFT acts similarly, integrating photoperiod and vernalization signals
LFY (LEAFY)	Controls floral meristem identity and flower development	DcLFY regulates floral meristem transition and organ formation
SOC1 (SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1)	Integrates environmental signals to activate FT	DcSOC1 has a similar function in regulating DcFT
FLC (FLOWERING LOCUS C)	Acts as a repressor of flowering until vernalization occurs	DcFLC prevents flowering before cold exposure suppresses its expression
GI (GIGANTEA)	Regulates circadian rhythms and flowering response to photoperiod	DcGI influences DcFT expression under long-day conditions

Role of Epigenetic Modifications in Flowering Regulation

Epigenetic modifications, which involve reversible changes to gene expression without altering DNA sequences, play an essential role in controlling flowering in carrots. These modifications help the plant remember environmental signals, such as cold exposure, ensuring that flowering occurs at the right time.

1. Histone Modifications

- Histones are proteins that help package DNA, and their modifications influence the accessibility of flowering genes.
- Before vernalization, floral repressors like *DcFLC* remain active due to specific histone methylation patterns, preventing premature flowering.
- Cold exposure triggers histone demethylation and acetylation, leading to the suppression of *DcFLC*, which allows flowering to proceed.

2. DNA Methylation

- DNA methylation is another mechanism that regulates flowering gene expression.
- In vernalized carrot plants, changes in DNA methylation patterns result in stable suppression of floral repressors, ensuring that flowering remains irreversible even after the cold period ends.
- This process, known as vernalization memory, ensures that once the plant transitions to reproductive growth, it does not revert to vegetative development.

3. Non-Coding RNAs

- Small RNAs and long non-coding RNAs (lncRNAs) also play a role in flowering control by interacting with flowering genes and modifying their expression.
- Some studies suggest that lncRNAs may regulate *DcFT* expression, adjusting flowering timing in response to environmental factors.

Epigenetic mechanisms provide an additional layer of regulation, ensuring that flowering occurs only under favourable conditions and preventing premature or delayed reproductive development.

Breeding Strategies for Improving Flowering Synchronization

Synchronizing flowering in carrot is a critical aspect of hybrid seed production. Since flowering is influenced by environmental factors like vernalization (cold exposure) and photoperiod (day length), breeding efforts focus on selecting genotypes with predictable and uniform flowering responses. Several strategies, including selection for vernalization and photoperiod responsiveness, hybridization, marker-assisted selection (MAS), and genetic engineering, help

improve flowering synchronization. These methods enable breeders to develop carrot varieties that flower consistently, increasing seed production efficiency.

Role of Hybridization and Marker-Assisted Selection (MAS)

Hybridization for Flowering Synchronization: Hybrid breeding is a widely used technique for improving flowering traits in carrots, particularly for hybrid seed production, where male-sterile and fertile pollinator lines must flower simultaneously.

- Crossing early- and late-flowering genotypes creates hybrids with more predictable flowering times.
- Recurrent selection and line breeding refine flowering uniformity over multiple generations.
- Cytoplasmic male sterility (CMS) systems, commonly used in carrot hybrid breeding, require careful synchronization of flowering between parental lines to optimize cross-pollination.

Marker-Assisted Selection (MAS) for Flowering Genes: Traditional selection for flowering traits can be slow due to environmental influences. Marker-assisted selection (MAS) accelerates the process by using molecular markers linked to genes regulating flowering time.

- Genetic markers associated with *DcFT*, *DcLFY*, and *DcFLC* genes allow breeders to identify plants with desirable flowering traits at an early stage.
- MAS enables breeders to predict flowering behavior without waiting for plants to mature, reducing the time required for field evaluations.
- Genomic selection (GS), an advanced form of MAS, analyzes the entire genome to predict flowering responses under different environmental conditions.

Potential of Genetic Engineering and CRISPR in Modifying Flowering Traits

Recent advancements in genetic engineering and CRISPR-Cas9 gene editing provide powerful tools for modifying flowering traits in a precise and controlled manner. These methods offer an alternative to traditional breeding, allowing for targeted improvements in flowering synchronization.

Genetic Engineering for Flowering Control

- Overexpression of *DcFT* (FLOWERING LOCUS T): Since *DcFT* promotes flowering, increasing its expression could reduce vernalization requirements, leading to earlier and more uniform flowering.
- Suppression of *DcFLC* (FLOWERING LOCUS C-like gene): *DcFLC* acts as a flowering repressor. Reducing its activity through genetic modification could accelerate flowering in genotypes that typically require long vernalization periods.
- Introduction of photoperiod-responsive genes from other species: Genes from photoperiod-sensitive plants could be introduced into carrots to fine-tune flowering times based on day length conditions.

CRISPR-Cas9 for Precision Breeding

CRISPR-Cas9 technology enables precise modifications of flowering-related genes, providing breeders with a faster and more efficient tool to improve flowering synchronization.

- Editing *DcFLC* to reduce its repression could lead to earlier and more consistent flowering, reducing dependency on long cold exposure.

- Modifying regulatory regions of *DcFT* could fine-tune its response to vernalization and photoperiod, ensuring uniform flowering under controlled conditions. Simultaneous editing of multiple flowering genes could allow breeders to optimize flowering behavior for specific climatic conditions.

References

1. Godwin, A., Pieralli, S., Sofkova-Bobcheva, S., & McGill, C. (2025). Natural genetic adaptation allows flexible reproductive behaviour: the case of wild carrot (Daucus carota L. subsp. carota) vs cultivated carrot (Daucus carota L. subsp. sativus). *Crop and Pasture Science*, *76*(1).
2. Linke, B., Alessandro, M. S., Galmarini, C. R., & Nothnagel, T. (2019). Carrot floral development and reproductive biology. *The carrot genome*, 27-57.
3. Spurr, C. J., & Lucas, A. M. (2020). Apiaceae seed production. In *Carrots and related Apiaceae crops* (pp. 18-34). Wallingford UK: CABI.
4. Verma, J., Singh, N., & Talwar, D. Color Spectrum of Carrot: Nutritional and Medicinal Properties.

Volume 3, Issue 2
February I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Soil Health and Conservative Ideas

ARTICLE ID: 0185

Ravneet Kaur

Bsc. (Hons.) Agriculture Student, PAU College of Agriculture, Ballowal Saunkhri

Soil is the fundamental concept of agriculture, which is essential for carrying out life processes of plants and animals. The concept soil health aims to protect soil from degradation caused by erosion (wind & water), deforestation, salinization, overgrazing by livestocks, excessive tillage practices etc. A healthy soil is essential for sustaining plant growth and animal life, regulates water, and supports nutrient cycle and protection against soil erosion. It represents the ability of soil to support healthy and sustainable environmental conditions to meet the demands of present as well as future generations. In this article we shall focus upon the different methods and conservative ideas for supporting soil health and fertility.

Importance of Maintaining Good Soil Health

"Soil Health" and "Soil Quality" are the terms that are used interchangeably. The characteristic of healthy soil include fertility, water holding capacity, optimum ph conditions, presence of organic matter, presence of active microorganisms, and accumulation of all essential nutrients. All these aspects contribute to growth of healthy plants. A good quality soil must be fertile enough so that it has all the essential nutrients such as nitrogen, potassium, phosphorus, calcium, magnesium accumulations in the soil. Plants may take these nutrients any time from soil, without their extra addition in soil. A good quality soil must maintain neutral ph around 6.5 to 8 for better adsorption of these ions as well to supports the microbial activities. Deceases in ph to less than 5 may lead to growth of fungi which is the major cause of fungal soil borne diseases. E.g. nematodes.

Innovative Approaches For The Conservation Of Soil Health

Soil health management systems aims to protect the soil structure and make the soil more productive for plant growth. These include

Conservation Tillage or Zero-Till: Excessive tillage may lead to change in soil structure and soil aggregates. It may expose the fertile soil layers to the surface which may wipe out easily due to erosion. It

may also leads to increased compactness of the soil thus reducing the macro-porosity and interfering plant growth. Reduced tillage practices may be beneficial for microbial activities in the undisturbed soil conditions and also help the farmers with increased profit due to low input of resources.

Use of Cover Crops: Cover crops play integral part in cropping system and provide soil cover in different seasons. These crops can be grown as annuals or biennials so that their crop residue can be used as mulching material for moisture retention. Growing of legume crops as cover crops helps in trapping excess nitrogen, and retains nitrate and other essential nutrients that are liable to leaching losses.

Organic Matter Management: Organic matter plays major role in integrating soil health. Organic matter is readily decomposed by micro-organisms and is the principle source of energy that acts as fuel in soil food web. It can be maintained, if the rate of addition of organic matter is greater than its utilization through decomposition. It can be maintained by incorporation of organic manures such as compost, green manures and animal wastes along with mulching.

Crop Rotation: Alteration of crops can lead to improvement of soil health and enhance the nutrient availability in the soil. Crop rotation helps in overcoming the soil borne disease cycles such as fungal diseases and also affects the life cycle of various pests. It also helps in management of weeds related to specific crops. Diversification of crops improves soil biodiversity and binding capacity of soil aggregates to plant roots thus preventing soil losses.

Nutrient management: Using diverse nutrient sources such as organic manures farm yard manure (FYM), poultry waste, bone meal, composts to soil with proper time and quantity management can improve the soil fertility and productivity. Maintaining proper soil ph improves the microbial activity and helps in improving the nutrient availability to plants as excess fertilizing may lead to toxic environmental conditions.

Reduced Pesticide Use: Excessive use of chemical fertilizers is the major cause of soil health deterioration. Overdose of nutrient fertilizers sprays leads to pest infestation which when controlled by chemical sprays infect the beneficial insects along with soil due to use of broad spectrum insecticides. These chemicals decrease the overall fertility of the soil.

Conclusion

Conserving soil health is essential for maintain agricultural productivity, ecological sustainability and overall planets health. Soil plays important role in regulating water systems, biodiversity, nutrient cycles, food production and mitigating climate changes. Therefore, challenges such as increasing population, urbanization, climate change, unstable agricultural practices are serious threats to the stability of soil. Implementing sustainable agricultural practices mentioned above may prevent the soil from further degradation. Protecting soil is not only agricultural issue but a combined global issue for sustaining life on Earth.

References

1. https://www.nrcs.usda.gov/conservation-basics/natural-resource-concerns/soils/soil-health

2. Introduction to Soils: Soil Quality by Lee Stivers in 202
3. Managing Soil Heath: Concepts and Practices by Charles White in 2017 (ARTICLE)