Volume 3, Issue 11
November I 2025

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Classification of Arthropod Pests Based on Feeding Habits

ARTICLE ID: 0284

Anas Khan

Student, School of Agriculture & Environmental Sciences, Shobhit Deemed-to-be University, Meerut-250110

Arthropods are the largest group of pests that damage agricultural crops and other plants. They feed on various plant parts such as leaves, stems, roots, flowers, and fruits, causing economic losses and reducing crop productivity. Classifying these pests based on their feeding habits helps in understanding their behavior, nature of damage, and in developing effective management strategies. This paper explains the main types of arthropod pests according to their feeding habits, along with common examples and suitable management practices.



1. Introduction

Arthropod pests occur in nearly all types of crops and environments. They include insects, mites, and other small organisms that damage plants by feeding on their tissues or transmitting diseases. Understanding the feeding behavior of these pests is crucial, as it determines the type of injury caused to plants and the control measures required.

Based on their feeding mechanism and mouthpart structure, arthropod pests are broadly classified into five categories:

1. Chewing pests
2. Piercing and sucking pests
3. Boring pests
4. Mining or scraping pests
5. Root feeders

Each group has unique characteristics, feeding patterns, and damage symptoms, which are discussed in detail below.

2. Classification Based on Feeding Habits

2.1 Chewing Pests

Chewing pests possess strong, well-developed mandibles and maxillae that allow them to bite and chew solid plant tissues. They feed on leaves, stems, flowers, or fruits, often leaving visible holes, irregular cuts, or defoliated portions. The extent of damage depends on the pest population and the crop stage.

Common examples: Caterpillars (*Spodoptera litura*, *Helicoverpa armigera*), grasshoppers, beetles, and weevils.

Damage symptoms

- Holes or irregular notches on leaves.
- Skeletonized leaves (only veins remain).
- Cut stems and defoliation in severe infestations.

Management Strategies

- Handpicking of larvae and egg masses.
- Application of neem-based or botanical insecticides.
- Spraying biological insecticides such as *Bacillus thuringiensis* (Bt).
- Encouraging natural enemies such as birds and predatory insects.
- Avoiding overuse of chemical insecticides to maintain ecological balance.

2.2 Piercing and Sucking Pests

These pests have needle-like mouthparts (stylets) that pierce plant tissues and suck out sap. This results in curling, yellowing, and stunted growth due to nutrient loss. They also serve as important vectors of viral and bacterial diseases.

Common examples: Aphids, whiteflies, jassids, leafhoppers, and mealybugs.

Damage Symptoms

- Yellowing and curling of leaves.
- Sticky honeydew secretion leading to sooty mold growth.
- Transmission of viral diseases such as leaf curl and mosaic.

Management Strategies

- Regular monitoring of pest population.
- Spraying neem oil or insecticidal soap.
- Introducing predators such as ladybird beetles and lacewings.
- Use of reflective mulches to repel whiteflies.
- Avoiding excessive nitrogen fertilizers which favor sap-sucking pests.

2.3 Boring Pests

Boring pests live and feed inside plant tissues by creating tunnels or galleries in stems, shoots, or fruits. As they feed internally, the damage is often hidden and difficult to detect until severe symptoms appear.

Common examples: Stem borers (in rice and maize), shoot borers (in brinjal and sugarcane), and fruit borers (in pomegranate and guava).

Damage symptoms

- Holes on stems, shoots, or fruits.
- Frass or sawdust-like material near entry holes.
- Wilting, drying, or breakage of shoots and branches.

Management Strategies

- Removal and destruction of infested plant parts.
- Installation of pheromone traps to attract and kill adults.
- Application of systemic insecticides at early crop stages.
- Field sanitation and crop rotation.
- Use of resistant crop varieties where available.

2.4 Mining and Scraping Pests

These pests feed on the internal or surface layers of leaves, creating characteristic mines, tunnels, or scraped patches. Leaf miners feed between the epidermal layers, while scraping pests remove the green chlorophyll layer, reducing photosynthetic efficiency.

Common examples: Leaf miners (*Liriomyza trifolii*), citrus leaf miners, and black-headed caterpillars.

Damage Symptoms

- Zigzag or serpentine mines visible on leaves.
- Brown or dry patches on leaf surfaces.
- Premature leaf fall and reduced photosynthesis.

Management Strategies

- Collection and destruction of affected leaves.
- Use of light or sticky traps to attract adult moths.
- Application of neem oil or spinosad formulations.
- Encouraging parasitoids like *Diglyphus isaea* for natural control.
- Avoiding excessive pesticide use that harms beneficial insects.

2.5 Root Feeders

Root feeders attack underground plant parts such as roots and rootlets, leading to poor nutrient uptake and stunted growth. The damage is often noticed only when plants show wilting or yellowing despite adequate irrigation and nutrition.

Common examples: White grubs, termites, and root aphids.

Damage symptoms

- Yellowing and wilting of plants.
- Easy uprooting due to damaged roots.
- Uneven plant growth and patchy field appearance.

Management Strategies

- Deep ploughing during summer to expose and kill larvae.
- Application of neem cake or *Trichoderma* in soil.
- Maintaining field hygiene and avoiding excess organic matter that attracts pests.
- Flooding fields where appropriate to control soil-dwelling stages.
- Using entomopathogenic nematodes and fungi as biological control agents.

3. Importance of Classification

Classifying arthropod pests based on feeding habits provides valuable insight into their biology, ecology, and mode of attack. It helps farmers, researchers, and extension workers identify pest damage accurately and choose effective management strategies. Furthermore, this classification serves as the foundation for Integrated Pest Management (IPM) programs, promoting sustainable and eco-friendly pest control.

4. Conclusion

Arthropod pests represent a major challenge in modern agriculture due to their diversity and adaptability. By understanding their feeding habits, farmers can predict the nature of damage and select appropriate control measures. Each pest category—chewing, sucking, boring, mining, or root feeding—requires specific management strategies. Integrating cultural, biological, and mechanical methods under the IPM framework minimizes pest losses while maintaining environmental safety.

References

1. Dhaliwal, G. S., & Arora, R. (2019). *Principles of Insect Pest Management.* Kalyani Publishers, India.
2. ICAR. (2022). *Integrated Pest Management Guidelines.* Indian Council of Agricultural Research, New Delhi.