Volume 4, Issue 11
November I 2026

Agri Roots
e- Magazine

www.agrirootsmagazine.in

ISSN: 2583-9071

Citronella: Cultivation and Essential Oil Production Technology

ARTICLE ID: 0445

Deepak Kumar

School of Agriculture and Environmental Science, Shobhit University, Meerut, Uttar Pradesh, India.

Citronella is an important aromatic and essential-oil-producing grass belonging to the family Poaceae. It is primarily cultivated for its fragrant essential oil, which is widely used in perfumes, cosmetics, soaps, detergents, candles and insect-repellent products. Citronella oil is also used in aromatherapy and certain pharmaceutical and personal-care preparations.

The two commercially important species are *Cymbopogon winterianus* (Java citronella) and *Cymbopogon nardus* (Ceylon citronella). Java citronella is generally preferred for commercial oil production because of its favourable oil yield and quality. The major constituents of citronella oil include citronellal, citronellol and geraniol, although their proportions vary with species, variety, growing conditions and distillation methods.



1. Botanical Description

Citronella is a perennial, tufted aromatic grass cultivated mainly for its leaves and aerial parts, which contain essential oil.

- **Common Name:** Citronella grass
- **Scientific Name:** *Cymbopogon winterianus* (Java citronella)
- **Family:** Poaceae
- **Plant Type:** Perennial aromatic grass
- **Propagation:** Slips or rooted clumps
- **Economic Part:** Leaves and aerial parts
- **Main Product:** Essential oil

2. Climate and Soil Requirements

2.1 Climate

Citronella grows well under warm and humid tropical and subtropical conditions. A temperature range of approximately 20–30°C is generally favourable for vigorous vegetative growth. Adequate sunlight supports healthy tillering, leaf development and essential oil production. The crop can tolerate short dry periods once established, but prolonged moisture stress may reduce growth and oil yield.

2.2 Soil

Citronella can be cultivated in a range of soil types; however, well-drained loamy to sandy-loam soils rich in organic matter are generally preferred. Good soil drainage is essential because prolonged waterlogging can damage the root system and adversely affect plant growth. Soil fertility and pH should be assessed before planting to support proper nutrient management.

3. Land Preparation

The field should be thoroughly prepared through ploughing and harrowing to obtain a fine, level and weed-free soil surface. Farmyard manure or well-decomposed compost may be incorporated during final land preparation to improve soil structure and fertility. Proper drainage arrangements should be made, particularly in areas receiving heavy rainfall.

4. Planting Material and Propagation

Citronella is commercially propagated through slips obtained by separating healthy and vigorous clumps from established mother plants. Healthy slips should possess a well-developed root system and sufficient basal growth to ensure successful establishment.

Before planting, excess leaves may be trimmed to reduce moisture loss and facilitate handling. Planting material should be selected from healthy, disease-free mother clumps to maintain crop vigour and uniformity.

5. Planting and Spacing

Healthy slips are planted in properly prepared fields at suitable spacing. A spacing of approximately 60 × 60 cm is commonly adopted, although the optimum spacing may vary depending on species, variety, soil fertility, rainfall and production system.

Planting should preferably be undertaken during the rainy season when adequate soil moisture is available for establishment. Irrigation should be provided when rainfall is insufficient.

6. Nutrient Management

Citronella responds well to balanced nutrient management because repeated harvesting of leaves and aerial biomass removes considerable quantities of nutrients from the soil.

Farmyard manure and well-decomposed compost can be incorporated to improve soil organic matter and nutrient availability. Nitrogen is particularly important for vegetative growth, tillering and leaf production, while phosphorus and potassium support root development and overall plant growth.

Nutrient application should be based on soil-test results and locally recommended practices. Split application of nitrogen may be beneficial because the crop produces repeated vegetative growth and is harvested several times during the growing season.

7. Irrigation Management

Citronella requires adequate soil moisture for continuous vegetative growth and regeneration after harvesting. Irrigation is particularly important during establishment, prolonged dry periods and after cutting. The frequency of irrigation depends on soil type, rainfall, temperature and crop growth stage. Light and timely irrigation is generally preferred over excessive watering. Proper drainage should be maintained to prevent waterlogging and root damage.

8. Weed Management

Weeds compete with citronella for nutrients, moisture, sunlight and space, particularly during the early stages of crop establishment. Effective weed management

helps ensure healthy plant growth and better field productivity.

Important weed-management practices include:

- Hand weeding during the early growth stages.
- Hoeing and inter-cultivation to control emerging weeds.
- Mulching to suppress weed growth and conserve soil moisture.
- Maintaining a clean field through regular monitoring.
- Using suitable herbicides only when recommended for citronella and applied according to approved label directions.

9. Pest and Disease Management

Citronella is comparatively hardy, but insect pests and diseases may occur under favourable environmental conditions. Their incidence can vary with the location, season and crop-management practices.

An Integrated Pest Management (IPM) approach should be followed. Important preventive measures include:

- Using healthy and disease-free planting material.
- Maintaining proper field drainage.
- Removing severely affected plant parts.
- Following field sanitation practices.
- Avoiding excessive and imbalanced fertilizer application.
- Regularly monitoring the crop for pest and disease symptoms.

Pest or disease-specific management measures should be adopted only after correct identification of the causal organism or pest.

10. Harvesting

The first harvest is generally taken several months after planting, once the crop has developed sufficient vegetative growth. Subsequent harvests can be obtained at regular intervals depending on climatic conditions, plant vigour and crop-management practices.

The leaves and aerial parts are cut above the ground using suitable harvesting tools, leaving sufficient basal portions for regrowth. Harvesting should be carried out at an appropriate stage to obtain good biomass and essential oil quality. The harvested material should be transported promptly for distillation to minimize deterioration and avoid unnecessary loss of volatile constituents.

11. Essential Oil Production Technology

Essential oil is primarily extracted from citronella grass through steam distillation or hydro-distillation. Steam distillation is widely used for commercial-scale extraction because it allows the recovery of volatile oil from plant material.

11.1 Preparation of Plant Material

Freshly harvested grass is cleaned to remove soil, weeds and other foreign materials. The plant material may be chopped into smaller pieces to facilitate uniform steam penetration. Proper handling and timely processing are important to minimize mechanical damage and the loss of volatile constituents.

11.2 Steam Distillation

The prepared grass is loaded into a distillation still. Steam is passed through the plant material, causing the volatile essential oil constituents to vaporize along with the steam. The vapour mixture is then directed

through a condenser, where it is cooled and converted into liquid form.

11.3 Condensation and Oil Separation

The condensed liquid contains essential oil and water, commonly known as distillation water or hydrosol. The essential oil is separated using a suitable oil separator based on differences in density and phase behaviour. The recovered oil is collected carefully to minimize contamination and handling losses.

11.4 Storage of Essential Oil

The extracted oil should be filtered, if necessary, and stored in clean, dry, airtight containers, preferably made of suitable glass or compatible material. Containers should be protected from direct sunlight, excessive heat and moisture to maintain oil quality during storage.

12. Chemical Constituents and Uses

Citronella essential oil contains several volatile compounds that contribute to its characteristic fragrance and commercial value. The major constituents include:

- **Citronellal:** An important fragrance compound used in perfumery and fragrance formulations.
- **Citronellol:** A fragrant alcohol used in perfumes, cosmetics and personal-care products.
- **Geraniol:** A commercially valuable aromatic compound used in fragrance and cosmetic industries.
- **Geranyl acetate:** An ester contributing to the characteristic fragrance of the oil.
- **Limonene:** A minor volatile constituent found in varying proportions.

The composition of citronella oil varies with species, variety, environmental conditions, harvesting stage and extraction method.

Major applications of citronella oil include:

1. Perfumes and fragrance formulations.
2. Soaps, detergents and household cleaning products.
3. Cosmetics and personal-care products.
4. Candles and aromatic preparations.
5. Insect-repellent formulations.
6. Aromatherapy and selected industrial applications.

13. Yield and Economic Importance

Citronella is a perennial aromatic crop capable of producing repeated harvests over several years under suitable management. Its productivity and essential oil yield depend on species, variety, planting density, soil fertility, climatic conditions, harvesting stage and distillation efficiency.

The crop offers opportunities for farm diversification because both biomass and essential oil have commercial applications. Establishment of small-scale distillation units, cooperative processing facilities and farmer-producer organizations can help improve access to processing and marketing infrastructure.

However, profitability depends on the cost of planting material, fertilizers, labour, irrigation, harvesting, distillation, transportation and market prices. Farmers should assess local demand, oil quality requirements and available processing facilities before undertaking commercial cultivation.

Conclusion

Citronella is an important aromatic crop with considerable potential for essential-oil production. Successful cultivation requires suitable planting

material, well-drained soil, balanced nutrient supply, adequate irrigation, effective weed management and timely harvesting. Steam distillation is an important technology for extracting citronella essential oil, which is widely used in fragrance, cosmetic, personal-care and insect-repellent industries.

With appropriate cultivation practices, efficient post-harvest handling and access to suitable distillation and marketing facilities, citronella can contribute to agricultural diversification, rural employment and additional income opportunities for farmers.

References

1. CSIR-CIMAP. *Cultivation Practices of Aromatic and Medicinal Plants*. Central Institute of Medicinal and Aromatic Plants, Lucknow, India.
2. Singh, A. K. (2004). *Medicinal and Aromatic Plants: Crops and Uses*. Agrobios, Jodhpur, India.
3. CSIR. *The Wealth of India: Raw Materials*. Council of Scientific and Industrial Research, New Delhi, India.
4. Farooqi, A. A., & Sreeramu, B. S. (2004). *Cultivation of Medicinal and Aromatic Crops*. Universities Press, India.
5. Board of Science and Technology for Rural Areas (BOSTID). *Aromatic Plants and Essential Oils*. National Research Council, Washington, D.C., USA.